

Dolor Literatur 99.1

Weiterführende Literatur (Literaturangaben von PD Dr. Ralf Baron)

1 Angel, I. F., H. J. Gould, Jr., M. E. Carey: Intrathecal morphine pump as a treatment option in chronic pain of nonmalignant origin. *Surg Neurol* 49 (1998) 92-98; discussion 98-99

2 Archer, A. G., V. C. Roberts, P. J. Watkins: Blood flow patterns in painful diabetic neuropathy. *Diabetologia* 27 (1984) 563-567

3 Attal, N., L. Brasseur, M. Chauvin et al.: A case of 'pure' dynamic mechano-allodynia due to a lesion of the spinal cord: pathophysiological considerations. *Pain* 75 (1998) 399-404

4 Attal, N., L. Brasseur, F. Parker et al.: Effects of Gabapentin on the Different Components of Peripheral and Central Neuropathic Pain Syndromes: A Pilot Study. *Eur Neurol* 40 (1998) 191-200

5 Backonja, M. M.: Reflex sympathetic dystrophy/sympathetically maintained pain/causalgia: the syndrome of neuropathic pain with dysautonomia. *Semin Neurol* 14 (1994) 263-271

6 Backonja, M., G. Arndt, K. A. Gombar et al.: Response of chronic neuropathic pain syndromes to ketamine: a preliminary study [published erratum appears in *Pain* 1994 Sep;58(3):433]. *Pain* 56 (1994) 51-57

7 Baron, R., M. Saguer: Postherpetic neuralgia. Are C-nociceptors involved in signalling and maintenance of tactile allodynia? *Brain* 116 (1993) 1477-1496

8 Baron, R., C. Maier: Painful neuropathy: C-nociceptor activity may not be necessary to maintain central mechanisms accounting for dynamic mechanical allodynia. *Clin J Pain* 11 (1995) 63-69

9 Benedetti, F., S. Vighetti, M. Amanzio et al.: Dose-response relationship of opioids in nociceptive and neuropathic postoperative pain. *Pain* 74 (1998) 205-211

10 Bennett, G. J., Y. K. Xie: A peripheral mononeuropathy in rat that produces disorders of pain sensation like those seen in man [see comments]. *Pain* 33 (1988) 87-107

11 Bennett, G. J., J. L. Ochoa: Thermographic observations on rats with experimental neuropathic pain. *Pain* 45 (1991) 61-67

12 Bennett, G. J.: An animal model of neuropathic pain: a review. *Muscle Nerve* 16 (1993) 1040-1048

13 Bennett, G. J.: Neuropathic Pain. In: Wall, P. D., Melzack, R. (eds.): *Textbook of Pain*, 3.ed. Churchill, Livingstone (1994) 201-224

14 Bennett, G. J.: Animal models of neuropathic pain. In: Gebhart, G. J., D. L. Hammond, T. S. Jensen (ed.): *Proceedings of the 7th World Congress on Pain*. Puplicher, Seattle (1994) 495-510

15 Bennett, G. J.: Neuropathic pain. In: Wall, P. D., R. Melzack (ed.): *Textbook of Pain*. Puplicher, Edinburgh (1994) 201-224

16 Bennett, D. L., J. French, J. V. Priestley et al.: NGF but not NT-3 or BDNF prevents the A fiber sprouting into lamina II of the spinal cord that occurs following axotomy. *Mol Cell Neurosci* 8 (1996) 211-220

17 Boureau, F., J. F. Doubrere, M. Luu: Study of verbal description in neuropathic pain. *Pain* 42 (1990) 145-152

- 18 Britland, S. T., R. J. Young, A. K. Sharma et al.: Acute and remitting painful diabetic polyneuropathy: a comparison of peripheral nerve fibre pathology. *Pain* 48 (1992) 361-370
- 19 Byas-Smith, M. G., M. B. Max, J. Muir et al.: Transdermal clonidine compared to placebo in painful diabetic neuropathy using a two-stage 'enriched enrollment' design. *Pain* 60 (1995) 267-274
- 20 Casale, R., M. Elam: Normal sympathetic nerve activity in a reflex sympathetic dystrophy with marked skin vasoconstriction. *J Auton Nerv Syst* 41 (1992) 215-219
- 21 Chabal, C., L. Jacobson, A. Mariano et al.: The use of oral mexiletine for the treatment of pain after peripheral nerve injury. *Anesthesiology* 76 (1992) 513-517
- 22 Chapman, V., R. Suzuki, H. L. Chamarette et al.: Effects of systemic carbamazepine and gabapentin on spinal neuronal responses in spinal nerve ligated rats. *Pain* 75 (1998) 261-272
- 23 Cherny, N. I., H. T. Thaler, H. Friedlander-Klar et al.: Opioid responsiveness of cancer pain syndromes caused by neuropathic or nociceptive mechanisms: a combined analysis of controlled, single-dose studies. *Neurology* 44 (1994) 857-861
- 24 Cho, H. J., D. S. Kim, D. S. Lee et al.: Changes in the alpha 2-adrenergic receptor subtypes gene expression in rat dorsal root ganglion in an experimental model of neuropathic pain. *Neuroreport* 29 (1997) 3119-3122
- 25 Choi, B., M. C. Rowbotham: Effect of adrenergic receptor activation on post-herpetic neuralgia pain and sensory disturbances. *Pain* 69 (1997) 55-63
- 26 Chung, K., B. H. Lee, Y. W. Yoon et al.: Sympathetic sprouting in the dorsal root ganglia of the injured peripheral nerve in a rat neuropathic pain model. *J Comp Neurol* 376 (1996) 241-252
- 27 Chung, K., Y. W. Yoon, J. M. Chung: Sprouting sympathetic fibers form synaptic varicosities in the dorsal root ganglion of the rat with neuropathic injury. *Brain Res* 751 (1997) 275-280
- 28 Ciccone, D. S., E. B. Bandilla, W. Wu: Psychological dysfunction in patients with reflex sympathetic dystrophy. *Pain* 71 (1997) 323-333
- 29 Culp, W. J., J. Ochoa, M. Cline et al.: Heat and mechanical hyperalgesia induced by capsaicin. Cross modality threshold modulation in human C nociceptors. *Brain* 112 (1989) 1317-1331
- 30 Daemen, M. A., H. A. Kurvers, P. J. Kitslaar et al.: Neurogenic inflammation in an animal model of neuropathic pain. *Neurol Res* 20 (1998) 41-45
- 31 DelleMijn, P. L., J. A. Vanneste: Randomised double-blind active-placebo-controlled crossover trial of intravenous fentanyl in neuropathic pain. *Lancet* 349 (1997) 753-758
- 32 Devulder, J. E., L. Ghys, W. Dhondt et al.: Neuropathic pain in a cancer patient responding to subcutaneously administered lignocaine. *Clin J Pain* 9 (1993) 220-223
- 33 Dickenson, A. H.: NMDA receptor antagonists: interactions with opioids. *Acta Anaesthesiol Scand* 41 (1997) 112-115
- 34 Dotson, R. M.: Clinical neurophysiology laboratory tests to assess the nociceptive system in humans. *J Clin Neurophysiol* 14 (1997) 32-45
- 35 Drummond, P. D., S. Skipworth, P. M. Finch: alpha 1-adrenoceptors in normal and hyperalgesic human skin. *Clin Sci (Colch)* 91 (1996) 73-77

- 36 Eija, K., T. Tiina, N. J. Pertti: Amitriptyline effectively relieves neuropathic pain following treatment of breast cancer. *Pain* 64 (1996) 293-302
- 37 Eisenberg, E., D. Pud: Can patients with chronic neuropathic pain be cured by acute administration of the NMDA receptor antagonist amantadine? *Pain* 74 (1998) 337-339
- 38 Ertas, M., A. Sagduyu, N. Arac et al.: Use of levodopa to relieve pain from painful symmetrical diabetic polyneuropathy. *Pain* 75 (1998) 257-259
- 39 Felsby, S., J. Nielsen, L. Arendt-Nielsen et al.: NMDA receptor blockade in chronic neuropathic pain: a comparison of ketamine and magnesium chloride. *Pain* 64 (1996) 283-291
- 40 Fields, H. L., M. C. Rowbotham: Multiple mechanisms of neuropathic pain: a clinical perspective. In: Gebhart, G. F., D. L. Hammond, T. S. Jensen (ed.): *Proceedings of the 7th World Congress on Pain*. Publisher, Seattle (1994) 437-454
- 41 Filshie, J.: The non-drug treatment of neuralgic and neuropathic pain of malignancy. *Cancer Surv* 7 (1988) 161-193
- 42 Fromm, G. H.: Baclofen as an adjuvant analgesic. *J Pain Symptom Manage* 9 (1994) 500-509
- 43 Galer, B. S.: Neuropathic pain of peripheral origin: advances in pharmacologic treatment. *Neurology* 45 (1995) S17-25; discussion S35-16
- 44 Galer, B. S., J. Harle, M. C. Rowbotham: Response to intravenous lidocaine infusion predicts subsequent response to oral mexiletine: a prospective study. *J Pain Symptom Manage* 12 (1996) 161-167
- 45 Galer, B. S., M. P. Jensen: Development and preliminary validation of a pain measure specific to neuropathic pain: the Neuropathic Pain Scale. *Neurology* 48 (1997) 332-338
- 46 Gillin, S., L. S. Sorkin: Gabapentin reverses the allodynia produced by the administration of anti-GD2 ganglioside, an immunotherapeutic drug. *Anesth Analg* 86 (1998) 111-116
- 47 Glazer, S., R. K. Portenoy: Systemic local anesthetics in pain control. *J Pain Symptom Manage* 6 (1991) 30-39
- 48 Gottrup, H., J. Nielsen, L. Arendt-Nielsen et al.: The relationship between sensory thresholds and mechanical hyperalgesia in nerve injury. *Pain* 75 (1998) 321-329
- 49 Gracely, R. H., S. A. Lynch, G. J. Bennett: Painful neuropathy: altered central processing maintained dynamically by peripheral input [published erratum appears in *Pain* 1993 Feb;52(2):251-3] [see comments]. *Pain* 51 (1992) 175-194
- 50 Hsieh, J. C., M. Belfrage, S. Stone-Elander et al.: Central representation of chronic ongoing neuropathic pain studied by positron emission tomography. *Pain* 63 (1995) 225-236
- 51 Hunter, J. C., K. R. Gogas, L. R. Hedley et al.: The effect of novel anti-epileptic drugs in rat experimental models of acute and chronic pain. *Eur J Pharmacol* 324 (1997) 153-160
- 52 Iadarola, M. J., M. B. Max, K. F. Berman et al.: Unilateral decrease in thalamic activity observed with positron emission tomography in patients with chronic neuropathic pain. *Pain* 63 (1995) 55-64
- 53 Iadarola, M. J., M. B. Max, K. F. Berman et al.: Unilateral decrease in thalamic activity observed with positron emission tomography in patients with chronic neuropathic pain. *Pain* 63 (1995) 55-64

- 54 Jett, M. F., J. McGuirk, D. Waligora et al.: The effects of mexiletine, desipramine and fluoxetine in rat models involving central sensitization. *Pain* 69 (1997) 161-169
- 55 Jevtovic-Todorovic, V., D. F. Wozniak, S. Powell et al.: Clonidine potentiates the neuropathic pain-relieving action of MK-801 while preventing its neurotoxic and hyperactivity side effects. *Brain Res* 781 (1998) 202-211
- 56 Jones, A.: The pain matrix and neuropathic pain [editorial]. *Brain* 121 (1998) 783-784
- 57 Jones, D. L., L. S. Sorkin: Systemic gabapentin and S(+)-3-isobutyl-gamma-aminobutyric acid block secondary hyperalgesia. *Brain Res* 810 (1998) 93-99
- 58 Kajander, K. C., S. Wakisaka, G. J. Bennett: Spontaneous discharge originates in the dorsal root ganglion at the onset of a painful peripheral neuropathy in the rat. *Neurosci Lett* 138 (1992) 225-228
- 59 Kajander, K. C., G. J. Bennett: Onset of a painful peripheral neuropathy in rat: a partial and differential deafferentation and spontaneous discharge in A beta and A delta primary afferent neurons. *J Neurophysiol* 68 (1992) 734-744
- 60 Kauppila, T., X. J. Xu, W. Yu et al.: Dextromethorphan potentiates the effect of morphine in rats with peripheral neuropathy [In Process Citation]. *Neuroreport* 9 (1998) 1071-1074
- 61 Kim, S. H., J. M. Chung: Sympathectomy alleviates mechanical allodynia in an experimental animal model for neuropathy in the rat. *Neurosci Lett* 134 (1991) 131-134
- 62 Kim, S. H., J. M. Chung: An experimental model for peripheral neuropathy produced by segmental spinal nerve ligation in the rat. *Pain* 50 (1992) 355-363
- 63 Kim, S. H., H. S. Na, K. Sheen et al.: Effects of sympathectomy on a rat model of peripheral neuropathy. *Pain* 55 (1993) 85-92
- 64 Kim, K. J., Y. W. Yoon, J. M. Chung: Comparison of three rodent neuropathic pain models. *Exp Brain Res* 113 (1997) 200-206
- 65 Kim, H. J., H. S. Na, B. Sung et al.: Amount of sympathetic sprouting in the dorsal root ganglia is not correlated to the level of sympathetic dependence of neuropathic pain in a rat model. *Neurosci Lett* 245 (1998) 21-24
- 66 Kingery, W. S.: A critical review of controlled clinical trials for peripheral neuropathic pain and complex regional pain syndromes. *Pain* 73 (1997) 123-139
- 67 Kishore-Kumar, R., S. C. Schafer, B. A. Lawlor et al.: Single doses of the serotonin agonists buspirone and m- chlorophenylpiperazine do not relieve neuropathic pain. *Pain* 37 (1989) 223-227
- 68 Kishore-Kumar, R., M. B. Max, S. C. Schafer et al.: Desipramine relieves postherpetic neuralgia. *Clin Pharmacol Ther* 47 (1990) 305-312
- 69 Klepstad, P., P. C. Borchgrevink: Four years' treatment with ketamine and a trial of dextromethorphan in a patient with severe post-herpetic neuralgia [see comments]. *Acta Anaesthesiol Scand* 41 (1997) 422-426
- 70 Kloke, M., K. Hoffken, H. Olbrich et al.: Anti-depressants and anti-convulsants for the treatment of neuropathic pain syndromes in cancer patients. *Onkologie* 14 (1991) 40-43
- 71 Koltzenburg, M., H. E. Torebjork, L. K. Wahren: Nociceptor modulated central sensitization causes mechanical hyperalgesia in acute chemogenic and chronic neuropathic pain. *Brain* 117 (1994) 579-591

- 72 Koltzenburg, M., H. E. Torebjörk, L. K. Wahren: Nociceptor modulated central sensitization causes mechanical hyperalgesia in acute chemogenic and chronic neuropathic pain. *Brain* 117 (1994) 579-591
- 73 Kontinen, V. K., S. Paananen, E. Kalso: The effects of the alpha2-adrenergic agonist, dexmedetomidine, in the spinal nerve ligation model of neuropathic pain in rats. *Anesth Analg* 86 (1998) 355-360
- 74 Kumar, K., C. Toth, R. K. Nath: Spinal cord stimulation for chronic pain in peripheral neuropathy. *Surg Neurol* 46 (1996) 363-369
- 75 Kurvers, H. A., G. J. Tangelder, J. G. De Mey et al.: Skin blood flow abnormalities in a rat model of neuropathic pain: result of decreased sympathetic vasoconstrictor outflow? *J Auton Nerv Syst* 63 (1997) 19-29
- 76 Lang, E., A. H. Hord, D. Denson: Venlafaxine hydrochloride (Effexor) relieves thermal hyperalgesia in rats with an experimental mononeuropathy. *Pain* 68 (1996) 151-155
- 77 Lanting, P., T. J. Faes, G. A. Ijff et al.: Autonomic and somatic peripheral nerve function and the correlation with neuropathic pain in diabetic patients. *J Neurol Sci* 94 (1989) 307-317
- 78 Lekan, H. A., K. Chung, Y. W. Yoon et al.: Loss of dorsal root ganglion cells concomitant with dorsal root axon sprouting following segmental nerve lesions. *Neuroscience* 81 (1997) 527-534
- 79 Levy, R. M., S. Lamb, J. E. Adams: Treatment of chronic pain by deep brain stimulation: long term follow-up and review of the literature. *Neurosurgery* 21 (1987) 885-893
- 80 Levy, R., J. Leiphart, C. Dills: Analgesic action of acute and chronic intraspinally administered opiate and alpha 2-adrenergic agonists in chronic neuropathic pain. *Stereotact Funct Neurosurg* 62 (1994) 279-289
- 81 Mao, J., D. J. Mayer, D. D. Price: Patterns of increased brain activity indicative of pain in a rat model of peripheral mononeuropathy. *J Neurosci* 13 (1993) 2689-2702
- 82 Mathisen, L. C., P. Skjelbred, L. A. Skoglund et al.: Effect of ketamine, an NMDA receptor inhibitor, in acute and chronic orofacial pain. *Pain* 61 (1995) 215-220
- 83 Max, M. B., R. Kishore-Kumar, S. C. Schafer et al.: Efficacy of desipramine in painful diabetic neuropathy: a placebo- controlled trial. *Pain* 45 (1991) p69; discussion 61-63
- 84 Max, M. B.: Treatment of post-herpetic neuralgia: antidepressants. *Ann Neurol* 35 Suppl (1994) S50-53
- 85 Max, M. B., M. G. Byas-Smith, R. H. Gracely et al.: Intravenous infusion of the NMDA antagonist, ketamine, in chronic posttraumatic pain with allodynia: a double-blind comparison to alfentanil and placebo. *Clin Neuropharmacol* 18 (1995) 360-368
- 86 McCaffery, M.: Gabapentin for lancinating neuropathic pain. *Am J Nurs* 98 (1998) 12
- 87 McCaffery, M.: Gabapentin for lancinating neuropathic pain. (1998)
- 88 McQuay, H. J.: Pharmacological treatment of neuralgic and neuropathic pain. *Cancer Surv* 7 (1988) 141-159
- 89 McQuay, H. J., D. Carroll, A. R. Jadad et al.: Dextromethorphan for the treatment of neuropathic pain: a double-blind randomised controlled crossover trial with integral n-of-1 design. *Pain* 59 (1994) 127-133

- 90 McQuay, H. J., M. Tramer, B. A. Nye et al.: A systematic review of antidepressants in neuropathic pain. *Pain* 68 (1996) 217-227
- 91 McQuay, H. J.: Opioid use in chronic pain. *Acta Anaesthesiol Scand* 41 (1997) 175-183
- 92 Mercadante, S., F. Lodi, M. Sapiro et al.: Long-term ketamine subcutaneous continuous infusion in neuropathic cancer pain. *J Pain Symptom Manage* 10 (1995) 564-568
- 93 Merren, M. D.: Gabapentin for treatment of pain and tremor: a large case series. *South Med J* 91 (1998) 739-744
- 94 Moriwaki, K., O. Yuge, H. Tanaka et al.: Neuropathic pain and prolonged regional inflammation as two distinct symptomatological components in complex regional pain syndrome with patchy osteoporosis--a pilot study. *Pain* 72 (1997) 277-282
- 95 Nelson, K. A., K. M. Park, E. Robinovitz et al.: High-dose oral dextromethorphan versus placebo in painful diabetic neuropathy and postherpetic neuralgia. *Neurology* 48 (1997) 1212-1218
- 96 Novakovic, S. D., E. Tzoumaka, J. G. McGivern et al.: Distribution of the tetrodotoxin-resistant sodium channel PN3 in rat sensory neurons in normal and neuropathic conditions. *J Neurosci* 18 (1998) 2174-2187
- 97 Nurmikko, T., A. Hietaharju: Effect of exposure to sauna heat on neuropathic and rheumatoid pain [published erratum appears in *Pain* 1992 Jun;49(3):419]. *Pain* 49 (1992) 43-51
- 98 Oaklander, A. L., K. Romans, S. Horasek et al.: Unilateral postherpetic neuralgia is associated with bilateral sensory neuron damage. *Ann Neurol* 44 (1998) 789-795
- 99 Ochoa, J. L., D. Yarnitsky: Mechanical hyperalgesias in neuropathic pain patients: dynamic and static subtypes. *Ann Neurol* 33 (1993) 465-472
- 100 Okuse, K., S. R. Chaplan, S. B. McMahon et al.: Regulation of expression of the sensory neuron-specific sodium channel SNS in inflammatory and neuropathic pain. *Mol Cell Neurosci* 10 (1997) 196-207
- 101 Omana-Zapata, I., M. A. Khabbaz, J. C. Hunter et al.: Tetrodotoxin inhibits neuropathic ectopic activity in neuromas, dorsal root ganglia and dorsal horn neurons. *Pain* 72 (1997) 41-49
- 102 Persson, J., G. Axelsson, R. G. Hallin et al.: Beneficial effects of ketamine in a chronic pain state with allodynia, possibly due to central sensitization. *Pain* 60 (1995) 217-222
- 103 Pfeifer, M. A., D. R. Ross, J. P. Schrage et al.: A highly successful and novel model for treatment of chronic painful diabetic peripheral neuropathy. *Diabetes Care* 16 (1993) 1103-1115
- 104 Portenoy, R. K., K. M. Foley, C. E. Inturrisi: The nature of opioid responsiveness and its implications for neuropathic pain: new hypotheses derived from studies of opioid infusions [see comments]. *Pain* 43 (1990) 273-286
- 105 Price, D. D., G. J. Bennett, A. Rafii: Psychophysical observations on patients with neuropathic pain relieved by a sympathetic block. *Pain* 36 (1989) 273-288
- 106 Pud, D., E. Eisenberg, A. Spitzer et al.: The NMDA receptor antagonist amantadine reduces surgical neuropathic pain in cancer patients: a double blind, randomized, placebo controlled trial. *Pain* 75 (1998) 349-354
- 107 Raja, S. N., Y. Choi, Y. Asano et al.: Arteriovenous differences in plasma concentrations of catechols in rats with neuropathic pain. *Anesthesiology* 83 (1995) 1000-1008

- 108 Raja, S. N.: Role of the sympathetic nervous system in acute pain and inflammation. *Ann Med* 27 (1995) 241-246
- 109 Rappaport, Z. H., M. Devor: Experimental pathophysiological correlates of clinical symptomatology in peripheral neuropathic pain syndromes. *Stereotact Funct Neurosurg* 54 (1990) 90-95
- 110 Ringkamp, M., S. Eschenfelder, E. J. Grethel et al.: Lumbar sympathectomy failed to reverse mechanical allodynia- and hyperalgesia-like behavior in rats with L5 spinal nerve injury. *Pain* 79 (1999) 143-153
- 111 Robbins, W. R., P. S. Staats, J. Levine et al.: Treatment of intractable pain with topical large-dose capsaicin: preliminary report. *Anesth Analg* 86 (1998) 579-583
- 112 Rosenberg, J. M., C. Harrell, H. Ristic et al.: The effect of gabapentin on neuropathic pain. *Clin J Pain* 13 (1997) 251-255
- 113 Rosner, H., L. Rubin, A. Kestenbaum: Gabapentin adjunctive therapy in neuropathic pain states. *Clin J Pain* 12 (1996) 56-58
- 114 Rowbotham, M. C., L. A. Reisner-Keller, H. L. Fields: Both intravenous lidocaine and morphine reduce the pain of postherpetic neuralgia. *Neurology* 41 (1991) 1024-1028
- 115 Rowbotham, M. C., P. S. Davies, H. L. Fields: Topical lidocaine gel relieves postherpetic neuralgia. *Ann Neurol* 37 (1995) 246-253
- 116 Rowbotham, M. C., P. S. Davies, C. Verkempinck et al.: Lidocaine patch: double-blind controlled study of a new treatment method for post-herpetic neuralgia. *Pain* 65 (1996) 39-44
- 117 Seltzer, Z., R. Dubner, Y. Shir: A novel behavioral model of neuropathic pain disorders produced in rats by partial sciatic nerve injury. *Pain* 43 (1990) 205-218
- 118 Seltzer, Z., S. Cohn, R. Ginzburg et al.: Modulation of neuropathic pain behavior in rats by spinal disinhibition and NMDA receptor blockade of injury discharge. *Pain* 45 (1991) 69-75
- 119 Shir, Y., Z. Seltzer: A-fibers mediate mechanical hyperesthesia and allodynia and C-fibers mediate thermal hyperalgesia in a new model of causalgiform pain disorders in rats. *Neurosci Lett* 115 (1990) 62-67
- 120 Shir, Y., A. Ratner, S. N. Raja et al.: Neuropathic pain following partial nerve injury in rats is suppressed by dietary soy. *Neurosci Lett* 240 (1998) 73-76
- 121 Siegan, J. B., A. T. Hama, J. Sagen: Suppression of neuropathic pain by a naturally-derived peptide with NMDA antagonist activity. *Brain Res* 755 (1997) 331-334
- 122 Stanton-Hicks, M., J. Salamon: Stimulation of the central and peripheral nervous system for the control of pain. *J Clin Neurophysiol* 14 (1997) 46-62
- 123 Stubhaug, A., H. Breivik: Long-term treatment of chronic neuropathic pain with the NMDA (N-methyl-D-aspartate) receptor antagonist ketamine [editorial; comment]. *Acta Anaesthesiol Scand* 41 (1997) 329-331
- 124 Study, R. E., M. G. Kral: Spontaneous action potential activity in isolated dorsal root ganglion neurons from rats with a painful neuropathy. *Pain* 65 (1996) 235-242
- 125 Takahashi, H., M. Miyazaki, T. Nanbu et al.: The NMDA-receptor antagonist ketamine abolishes neuropathic pain after epidural administration in a clinical case. *Pain* 75 (1998) 391-394

- 126 Tal, M., G. J. Bennett: Neuropathic pain sensations are differentially sensitive to dextropropofol. *Neuroreport* 5 (1994) 1438-1440
- 127 Tanelian, D. L., M. B. MacIver: Analgesic concentrations of lidocaine suppress tonic A-delta and C fiber discharges produced by acute injury. *Anesthesiology* 74 (1991) 934-936
- 128 Tasker, R. R., O. Vilela Filho: Deep brain stimulation for neuropathic pain. *Stereotact Funct Neurosurg* 65 (1995) 122-124
- 129 Tesfaye, S., J. Watt, S. J. Benbow et al.: Electrical spinal-cord stimulation for painful diabetic peripheral neuropathy [see comments]. *Lancet* 348 (1996) 1698-1701
- 130 Torebjörk, E., L. Wahren, G. Wallin et al.: Noradrenaline-evoked pain in neuralgia [see comments]. *Pain* 63 (1995) 11-20
- 131 Torebjörk, E., L. Wahren, G. Wallin et al.: Noradrenaline-evoked pain in neuralgia [see comments]. *Pain* 63 (1995) 11-20
- 132 Treede, R. D., K. D. Davis, J. N. Campbell et al.: The plasticity of cutaneous hyperalgesia during sympathetic ganglion blockade in patients with neuropathic pain. *Brain* 115 (1992) 607-621
- 133 Verdugo, R. J., J. L. Ochoa: Use and misuse of conventional electrodiagnosis, quantitative sensory testing, thermography, and nerve blocks in the evaluation of painful neuropathic syndromes. *Muscle Nerve* 16 (1993) 1056-1062
- 134 Verdugo, R. J., J. L. Ochoa: 'Sympathetically maintained pain.' I. Phentolamine block questions the concept [see comments]. *Neurology* 44 (1994) 1003-1010
- 135 Wakisaka, S., K. C. Kajander, G. J. Bennett: Abnormal skin temperature and abnormal sympathetic vasomotor innervation in an experimental painful peripheral neuropathy. *Pain* 46 (1991) 299-313
- 136 Watson, C. P., N. Babul: Efficacy of oxycodone in neuropathic pain: a randomized trial in postherpetic neuralgia [In Process Citation]. *Neurology* 50 (1998) 1837-1841
- 137 Yoon, Y. W., H. S. Na, J. M. Chung: Contributions of injured and intact afferents to neuropathic pain in an experimental rat model. *Pain* 64 (1996) 27-36
- 138 Zochodne, D. W.: Epineurial peptides: a role in neuropathic pain? *Can J Neurol Sci* 20 (1993) 69-72